

Effect of Cupressus Arizonica Essential Oil on Two Spotted Spider Mite (Tetranychus urticae) on Alii Fig (Ficus maclellandii)

Alireza Hasibi

Landscape Department of Tehran Municipality, District 12. Tehran, Iran.

Abstract: Two spotted spider mite (Tetranychus urticae) is one of the most widespread and serious pests of farm and greenhouse plants. Nowadays, the use of plant essentials is considered due to the harmful effects of chemical pesticides. This study was conducted to assess the toxic potential of Arizona cypress (Cupressus arizonica) essential oil as an effective acaricide on Tetranychus urticae on Alii Fig (Ficus maclellandii). This study was conducted using randomized complete design with 8 treatments (7 treatments and 1 control). The amount of dead mites and burned Alii Fig leaves were estimated after 24 hours. LD50 value was measured. The results showed that increasing essential oil concentrations increased the toxic effect of the essential. At the concentration of 2.5 ml* L⁻¹, the essential caused the highest level of mortality of mites. LD50 value was estimated 1.25 ml* L⁻¹. There was no burn on Alii Fig leaves up to concentration of 5 ml* L⁻¹ of the essential.

Keywords: Alii Fig, mite, plant essential oil, Arizona cypress

INTRODUCTION

Alii Fig (*Ficus maclellandii*) from the Moraceae family, is a widely used as an ornamental leaf plant in Iran. *Ficus maclellandii* is a species of fig plant native to India, Southeast Asia and China. It is an evergreen often grown as a houseplant in temperate climates. The leaves are 8–13 cm and uniquely dimorphic; with narrow leaves on the lower, sterile branches and broader leaves on the higher branches.

The use of herbal ingredients in pest control of house plants has positive impacts on environment and people health.

The two spotted spider mite (*Tetranychus urticae*), is one of the most damaging pests in agricultural, horticultural and greenhouse systems. It causes the reduce of chlorophyll content and photosynthesis in the leaf and eventually lead to leaf loss and plant weakness (Gorman *et al.*, 2001).

Large quantities of chemical pesticides are used every year to control the pest, which due to its short shelf life and high reproduction, it is resistant to pesticides, and practically repeated use of chemical pesticides only causes environmental pollution. And the negative effects on non-target beings. (Isman, 1999).Therefore, the use of herbal compounds such as essential oils and extracts has been considered as an alternative method of mite management especially in greenhouse conditions. (Cavalcanti *et al.*, 2010)

The essential oils of plants are well-known natural ingredients in pests control, as well as the repellent effects of plant essential oils such as peppermint on two-point spider mites (Moayeri *et al.*,

Spec. J. Agric. Sci, 2019, Vol, 5 (4): 10-14

2013).The essential oils are easy to extract and decompose quickly after use and adapt to the environment. In addition, most essential oils are not toxic to mammals, fish and birds and do not persist in soil and water (Ziaie *et al.*, 2015).

Arizona cypress (*Cupressus arizonica*) is an evergreen and resistant plant that tolerates a wide range of ecological conditions (Emami *et al.*, 2004) and is abundantly planted on the surface of the landscape. It was found that the essential oil of Arizona cypress has 46 compounds. Limonene (14.44%), ambulolone (13.25%) and alpha-pinene (11%) are the major constituents of this essential oil (Sedaghat et al., 2011).

Since the cypress trees are pruned every year in urban landscape and their waste is discarded, using this inexpensive and affordable source of the production of pesticides is economical and rational, so this experiment was conducted to evaluate the effect of Arizona cypress essence on two spotted spider mite on Alii Fig plant.

Materials and Methods

Branches and leaves of Arizona cypress were collected from the landscape of Park-e-Shahr of Tehran and transferred to the laboratory of Landscape Research, Education and Counseling Center of District 12 of Tehran Municipality, during autumn 2018 and dried at room temperature.

The essential oil of the plant was prepared by Clevenger apparatus and water and steam distillation from 100 g of cypress leaves in one liter of distilled water for 2 h. The essential oil was stored in dark glass at 4 ° C until testing. To prepare different concentrations of essential oil, stock solution was prepared from 2 cc of essential oil and twin 20 as a 2% emulsion, distilled water to 100 ml volume was added and other concentrations were prepared using different amounts of distilled water.

For the purpose of bioassay of mites, mite infected leaves of Alii Fig were prepared from greenhouse of Landscape Research, Education and Counseling Center of District 12 of Tehran Municipality and transferred to laboratory. The number of mature mites and eggs were counted on infected leaves.

Leaf immersion method was used to determine the lethal concentrations of different the essential oil concentrations. The infected leaves were immersed in essential oil solution for 5 seconds at concentrations of 0.15, 0.3, 0.6, 1.25, 2.5, 5 and 10 ml* L \cdot 1. The control treatment was considered distilled water. Treated leaves were exposed to air at room temperature to dry and transferred to 9-cm petri dishes with a layer of wet filter paper. The Petri cap was ventilated with a 3 cm diameter pore covered with mesh. Petri dishes were kept at 24 ± 2 ° C with 14 h light. The status of mites was assessed 24 hours later with binoculars and the number of killed mites was counted.

In order to evaluate the degree of burnout of leaves, they were immersed in various concentrations of the essential oil for 5 seconds and then exposed to air at room temperature for 20 to 30 minutes to dry and placed in Petri dish containing a layer of filter paper with a pore diameter of 3 cm.

After 24 hours, leaf evaluation was performed and scoring was 0-10, score 0 means no burns and score 10 indicates complete necrosis of leaf tissue.

The experiment was conducted in a completely randomized design with three replications. Data were analyzed using Minitab 16 software and Sigmaplot 11 software.

Results

The results indicate the effect of Arizona cypress essential oil on population decline of two spotted spider mite on Alii Fig leaf.

The mortality rate of mites at different concentrations of essential oil was significantly different $(p \le 0.01)$.

The percentage of mortality at a concentration of 0.3 ml* L ⁻¹ was only 1.75% and increased to 0.6 ml* L ⁻¹ at 27.5% and resulted in complete death and 100% pest at concentrations of 2.5 ml* L ⁻¹ and above (Figure 1).



Figure 1. Effect of Arizona cypress essential oil on mortality rate of two spotted spider mite in Alii Fig plant

The insecticidal properties of silver cedar essential oil have been proven in several cases the fumigant effect of essential oils of Cumin, Anise, Oregano and Eucalyptus on *Aphis gossipii* was confirmed (Tunc *et al.*, 1998). Also, the effect of essential oil of three rosemary, peppermint and savory herbs on two spotted spider mite was investigated and confirmed (Moayeri *et al.*, 2013).

Trend changes in mortality rate of different essential oil concentrations by a 3-parameter sigmoidal model [Mortality (%)=100/(1=exp(-(x-1.25)/0.39))]. LD10, LD50 and LD90 values were 0.4, 1.25 and 2.07 ml* L -1, respectively.

The insecticidal properties of silver cedar essential oil have been proven in several cases (Habibi *et al.*, 2012; Kabiri *et al.*, 2014; Heidari Zadeh *et al.*, 2012). The fumigant effect of essential oils of Cumin, Anise, Oregano and Eucalyptus on *Aphis gossipii* was confirmed (Tunc *et al.*, 1998). Also, the effect of essential oil of three rosemary, peppermint and savory herbs on two spotted spider mite was investigated and confirmed (Moayeri *et al.*, 2013).

Trend changes in mortality rate of different essential oil concentrations by a 3-parameter sigmoidal model [Mortality (%)=100/(1=exp(-(x-1.25)/0.39))]. LD10, LD50 and LD90 values were 0.4, 1.25 and 2.07 ml* L -1, respectively.



Figure 2. Changes in mortality rate of two spotted spider mite in Benjamin plant at different concentrations of Arizona cypress essential oil

in previous studies, LD50 content of ethanolic extract of bitter olive seed on peach green aphid was calculated to be 15 µl * ml-1, after 24 h (Salari *et al.*, 2011). Evaluation of toxicity of essential oil of some plants of mint family on two spotted spider mite revealed that thyme and peppermint essential oil with the LD 50 of 419 and 425 mg *l⁻¹ had the highest toxicity on two spotted spider mite, respectively (Kaveh *et al.*, 2014).

The results of this study showed that silver cedar extract at concentrations less than 5 ml^{*} L ⁻¹ did not cause any burns on Benjamin leaves. At concentrations of 5 and 10 ml^{*} L ⁻¹, 11 and 23% of burns were observed, respectively.

In the study of toxicity, the essential oils of 10 specious of mint family on two spotted spider mite at concentrations below 1600 ppm, the essential oils did not cause any burns on the leaves of bean plant (Kaveh *et al.*, 2014).

Due to complete death of ticks at concentrations of 2.5 to 10 ml* L -1 and no signs of burns on Benjamin leaves at 2.5 ml* L -1 and slight burn at 5 ml* L -1, the concentration range 2.5 to 5 ml* L -1 is recommended for the preparation of the pesticide. Essential oils of barley, artemisia and geranium had low germination effects at 12 ppm concentration in greenhouse cucumber (Yarahmadi *et al.*, 2012).

The findings of the present study indicate that Arizona cypress essential oil is can be used as a low risk pesticide to control mites on the Alii Fig plant. Due to the less destructive effects of plant compounds on humans and the environment, the essential oil of this plant is recommended for the manufacture of natural acaricides.

References

 Cavalcanti, S. C. H., Niculau, E. D. S., Blank, A. F., Câmara, C. A. G., Araújo, I. N., & Alves, P. B. (2010). Composition and acaricidal activity of Lippia sidoides essential oil against twospotted spider mite (Tetranychus urticae Koch). *Bioresource Technology*, 101(2), 829-832. Spec. J. Agric. Sci, 2019, Vol, 5 (4): 10-14

- Gorman, K., Hewitt, F., Denholm, I., & Devine, G. J. (2002). New developments in insecticide resistance in the glasshouse whitefly (Trialeurodes vaporariorum) and the two-spotted spider mite (Tetranychus urticae) in the UK. *Pest Management Science: formerly Pesticide Science*, 58(2), 123-130.
- Habibi, F., Moarrefi, M. & Rafii, Z., (2012). Fumigant toxicity and repellency effect of essential oils of Pinus eldarica and Cupressus arizonica on adults of flour moth, Ephestia kuehniella Zeller. (Lep., Pyralidae). *Journal of Entomological Research*, 4, 215-225.
- Hassanzadeh Khayyat, M., Emami, S. A., Rahimizadeh, M., Fazly-Bazzaz, B. S., & Assili, J. (2005). Chemical constituents of Cupressus sempervirens L. cv. Cereiformis Rehd. essential oils. *Iranian Journal of Pharmaceutical Sciences*, 1(1), 39-42.
- 5. Heidari Zadeh, M., Lotfi, V. & Ghanei, M. (2018). Evaluation of chemical compounds, antibacterial and allelopathic properties of cedar leaf extract (Cupressus arizonica). *Iran biology magazine*, 31.
- Isman, M. (1999). Pesticides based on plant essential oils. *Pesticide Outlook (United Kingdom)*. 68 – 72.
- 7. Kabiri, M., Mohammadi Sharif, M. & Kabirinasab, M. (2014). Biological effects of Arizona cypress, Cupressus arizonica against rice weevil, Sitophilus oryzae and sawtoothed grain beetle, Oryzaephilus surinamensis, *Journal of Plant Protection*, *37*, 11-21.
- Kaveh, M., Poorjavad, N. & Khaje Ali, J. (2014). Evaluation of contact toxicity of ten essential oils from Lamiaceae plants against Tetranychus urticae Koch (Acari: Tetranychidae), *Plant Pests Research*, 4, 39-49.
- 9. Moayeri, H., Pirayeshfar, F. & Kavousi, A. (2013). The repellent effect of three herbal essential oils on Tetranychus urticae, *Iranian Journal of Plant Protection Knowledge*, 44, 103-112.
- 10. Salari, A., Ahmadi, K. & Zamani, R. (2011). The effect of ethanolic extract of bitter olive (Melia azedarach L.) seed on two aphid species, *Journal of Herbal Medicines, 9*, 223-228.
- Sedaghat, MM, Sanei Dehkordi, A., Khanavi, M., Abai, M.R., Mohtarami, F. & Vatandoost, H. (2011). Chemical composition and larvicidal activity of essential oil of Cupressus arizonica E.L. Greene against malaria vector Anopheles stephensi Liston (Diptera: Culicidae), *Pharmacognosy Research, 3*, 135-139.
- 12. Tunc, I., & Sahinkaya, S. (1998). Sensitivity of two greenhous pests to vapors of essential oils, *Entomologia Experimentalis et Applicata, 86*, 183-187.
- Yarahmadi, F. and Rajabpoor, A. (2012). Efficacy of the essential oil of geranium in control of Bemisia tabaci Gen in greenhouse cucumbers, National Conference on Natural Products and Medicinal Plants, Iran.
- 14. Ziaie, M. & Hamzevi, F. (2015). Essential oils in integrated pest management programs, Journal of Agricultural and Natural Resources Engineering, 47, 26-30.