



# Risk Factors Associated with Urinary Tract Parasitic Infections among Students of Rivers State College of Health Science and Tecnology, Port Harcourt, Rivers State, Nigeria

Lebari Barine Gboeloh<sup>1\*</sup>, Nelly Onyinyechi Elechi<sup>2</sup>, Ike-Ihunwo Chikaire Ndamzi<sup>2</sup>

<sup>1</sup> Department of Biology, Faculty of Natural and Applied Sciences, Ignatius Ajuru University of Education, Port Harcourt, Nigeria

<sup>2</sup> Department of Biomedical Technology, School of Biomedical Engineering Technology, Rivers State College of Health Science and Tecnology, Port Harcourt, Nigeria

## \*Corresponding Author

**Abstract:** Urinary tract infections are common non-viral diseases mostly transmitted through sexual activities. Our study investigated the prevalence and risk factors associated with these infections among students of Rivers State college of Health Sciences and Tecnology, Port Harcourt, Rivers State. A cross sectional study was conducted in which urine specimens from 353 students (89males, 253females) were collected. The samples were prepared for microscopic examination using wet mount method for observation of motile organisms and sedimentation technique for observation of eggs and cysts. Out of the 353 samples investigated, 107(30.3%) were infected. The parasites identified included *Trichomonas vaginalis*, *Schistosoma haematobium* and *Enterobius Vermicularis*. Of all the parasite species identified, *T. vaginalis* showed a statistically significant prevalence ( $p<0.05$ ). Although there was no significance relationship between age and prevalence of the parasites, the most infected age group was 21-25yrs (29%) followed by 16-20yrs (27%) while the least infection was found among students above the age of 40years (6%). More females (86.9%) were infected than males (13.1%) and this was statistically significant ( $p<0.05$ ). Similarly, more single women (92.5%) were infected than married women (7.5%). There was no significance relationship ( $p>0.05$ ) between nature of toilet used and prevalence of the parasites. The students that used bucket/ semi water system at home, water closet system at home and pit toilet at home had 54.2%, 25.5% and 22.4% respectively. The high prevalence rate observed in the study is a source of great concern. Health education, regular screening and treatment of infected persons will reduce the trend.

**Keywords:** Urinary Tract Infections, Risk factor, Prevalence, *T. vaginalis*, *E. vemicularis*, *S. haematobium*

## INTRODUCTION

*Trichomonas vaginalis* is a flagellate protozoan. It is one of the few parasites that inhabit the urogenital system of both males and females (Jatau *et al.*, 2006) and it has been isolated from nearly all parts of the

urogenital system including the uterus of infected individuals (Ijeoma *et al.*, 2018). Trichomoniasis, the disease caused by *T. vaginalis* is a pluralistic disease (Akinbo *et al.*, 2017) mostly transmitted through unprotected sexual intercourse and it is the third most common cause of vaginal symptoms behind bacterial vaginosis and candidiasis (Toth, 2013). It could also be transmitted non-sexually through contact with fomites and surfaces harbouring fluids from infected individuals (Nester *et al.*, 2001; Ijeoma *et al.*, 2018) including sharing of pants and other underwears with infected persons (Akinbo *et al.*, 2017).

The disease affects an estimated 180 million people worldwide (Bowden and Garnett, 2000; Uproft and Uproft, 2001). WHO (2012) reported that globally, an estimated 276.4 million new cases of the infection are recorded annually. In America alone, an estimated 3million people are infected yearly (WHO, 2001) and in Africa, an estimated 42.8million people are at risk of the infection (de Waaiji *et al.*, 2017; WHO, 2012). According to WHO (2002), women are more vulnerable to trichomoniasis than men, prevalence rate of 41.4%, 34%, 10.7%, 10.2% and 9.9% have been recorded among pregnant women in South Africa, Tanzania, Gabon, Congo and Central African Republic respectively.

In Nigeria, although the record is scanty, various degree of prevalence of the infection have been recorded in different part of the country (Ijeoma *et al.*, 2018; Ijase *et al.*, 2018, Olusola *et al.*, 2010; Akinbo *et al.*, 2017, Sam-Wobo *et al.*, 2012).

Although the disease is one of the commonest treatable sexually transmitted infections affecting public health (Hirt *et al.*, 2011; von Glehn *et al.*, 2017), it could cause secondary diseases such as pelvic inflammatory disease in women (Moodley *et al.*, 2002), lower birth weight in children of infected mothers (Cotch *et al.*, 1997), defective and low sperm count in men (Ryu *et al.*, 2015) and preterm labour (Mann *et al.*, 2010). Trichomoniasis has also been implicated as one of the high-risk factors in aggravation of HIV infection and transmission (Van de pol, 2007; Kissenger, 2013). Trichomoniasis is reportedly implicated in reduced immune activation, replication and cytokine production resulting increased viral load of people living with HIV (Quinlivan *et al.*, 2012). Other health implication of trichomoniasis include urethritis, vaginitis, cervicitis, prostatitis particularly in HIV infected individuals (Mahmoud *et al.*, 2015) and infertility (Fichorava, 2009; Goodkin *et al.*, 2005).

In most cases, the disease is asymptomatic in both males and females (Fouts and Kraus, 1980), hence it is not usually and easily diagnosed (Akinbo *et al.*, 2017). However, women are more vulnerable than men and an estimated 180 million females are infected globally (Swygard *et al.*, 2004).

## **Materials and Methods**

### **Study Area**

Rivers State College of Health Sciences and Technology is one of the tertiary institutions in Rivers State. It is sited in Obio/Akpor Local Government Area and lies on latitude 4<sup>o</sup>49'N and longitude 6<sup>o</sup>59'E. The college share the same climatic conditions similar to establishments within Obio/Akpor Local Government Area. The Local Government Area is characterised by unpredictable rainfall throughout the year with intermittent dry season.

### **Study Design and Study Population**

The study was a cross sectional research carried out among students of the college. The study was conducted within July, 2018 and January, 2019. The study population included admitted and registered students of the school residence in the hostel accommodation provided by the school. Official record from the registrar's office indicated that the population of students in the school was 3000 as at the time of the study.

### **Sample Size**

The sample size for this study was 353 students. This was determined using the method of Taro (1967):  $n=N/(1+(e)^2)$ .

Where: n = sample size

$N$  = population of study group (3000)

$e$  = margin error (0.05)

$n = N/(1+(e)^2)$

$n = 3000/(1+3000(0.05)^2)$

$n = 3000/ (1+ 3000 \times 0.0025)$

$n = 3000/ (1+7.5)$

$n = 3000 / 8.5$

$n = 353$

**Questionnaire:** A simple self-structured questionnaire was administered to all the sampled participants. The questionnaire was used to collect demographic data on age, sex, marital status and nature of toilet facility used.

### **Sample Collection**

Well labelled sterile sample bottles were distributed to the students and were allowed to go home with it having been instructed to collect their mid-stream early morning urine. They were advised to return the bottles within few hours after collection.

### **Laboratory Analysis**

The samples were transported to the research laboratory, Department of Biology, Ignatius Ajuru University of Education, Port Harcourt for parasitological examination.

**Wet mount test:** Wet mounts preparation was made by placing few drops of the urine on a sterile and clean glass slide, covered with a cover slip and examined under the microscope for motile flagellates of *Trichomonas vaginalis* using the X10 and X40 objectives.

**Sedimentation method:** To enhance accuracy, about 5mls of each of the fresh urine was mixed with equal volume of physiological saline. The mixture was properly shaken for harmonization and centrifuged at 3500rpm for 5minutes. The supernatant was discarded leaving the sediment at the bottom of the tube. A drop of the sediment was placed on a clean glass slide, covered with a cover slip and observed under the microscope for *T. vaginalis*.

### **Data Analysis**

Data generated was analyzed using One way analysis of variance (ANOVA) to determine the significant difference between variables at 0.05 significant level of prevalence.


### **Ethical Clearance**

The ethical clearance for this study was obtained from the Ethical Committee and Dean of Student Affairs of the college. The consent of the subjects was also sought and obtained.

## **Results**

The overall prevalence of urinary tract parasitic infection among the study group indicated that out of the 353 students examined, 107 (30.3%) were positive for at least one urinary tract parasitic disease (Fig.1). The urinary tract parasites isolated and identified from urine samples of the students investigated included *Trichomonas vaginalis* (Plate 1), Eggs of *Schistosoma haematobium* (Plate 2) and Egg of *Enteriobius Vermicularis* (Plate 3). The results indicated a statistically significance ( $P < 0.05$ ) prevalence of the infection among the students investigated. Out of the 107 infected students, 27(25.3%), 31(29.0%), 21(19.6%), 15(14.0), 7(6.5%) and 6(5.6%) belonged to the age range of 16-20yrs, 21-25yrs, 26-30yrs, 31-35yrs, 36-40yrs and 41yrs and above respectively (Table 1). The most infected age group was 21-25yrs (29%) followed by 16-20yrs while the least infection was found among students above the age of 40years. Age has no statistically significant ( $p > 0.05$ ) impact on the prevalence of the infections. More females 93(86.9%) were infected than males 14(13.1%). This shows a significance difference ( $p < 0.05$ ) suggesting that sex is an important factor in the prevalence of the infections.

The students that used bucket/ semi water system at home were more infected 58(54.2%) than those that use water closet system at home and in school 27(25.5%) while the students that use pit toilet at home had 24(22.4%). No significance difference ( $p>0.05$ ) exists in the prevalence of urinary parasitic infections in relation to the types of toilet system used.



**Figure 1:** overall prevalence of urinary tract parasitic infection

**Table 1:** Risk factors associated with urinary tract parasitic infections among students of Rivers State College of Health Sciences and Technology.

Risk Factors	No. Examined	No. Infected						
<b>Species of Parasites</b>								
AGE (yr)		T. <i>Vaginalis</i>	S. <i>haematobium</i>	E. <i>vemicularis</i>	Total (%)	Df	F	Sig.
16 - 20	97	25	0	2	27(25.2)	5	.597	.703
21 - 25	107	29	1	1	31(29.0)			
26 - 30	63	20	1	0	21 (19.6)			
31- 35	48	15	0	0	15 (14.0)			
36-40	26	3	3	1	7(5.6)			
Above 41	12	5	1	0	6(5.6)			
<b>Sex</b>								
Male	89	11	3	0	14(13.1)	1	12.328	.001
Female	253	86	3	4	93(86.9)			
<b>Marital Status</b>								
Single	256	89	6	4	99(92.5)	1	33.524	.000
Married	97	8	0	0	8(7.5)			
<b>Nature Of Toilet</b>								
Bucket system at home	353	52	3	3	58(54.2)	2	.033	.967
Water closet at school	353	25	1	1	27(25.5)			
Pit toilet at home	353	20	2	0	24(22.4)			
<b>Total</b>	<b>353</b>							

**Discussion**

Urinary tract parasitic infections have gradually become a major health concern among primary health practitioners especially in developing countries of the world. The infections have been implicated in the

emergence of several secondary diseases associated with the productive capacity of both males and females (Swygard *et al.*, 2004; Patel *et al.*, 2018).

In this study, the prevalence and risk factors associated with urinary tract parasitic infections among the study group were investigated. The parasites isolated and identified included *T. vaginalis*, *E. vermicularis* and *S. haematobium*.

The high prevalence rate (30.3%) recorded in our study is of great concern. *T. vaginalis* recorded the highest prevalence of 90.7% compared to the prevalence of other parasites identified. This was also significantly higher than other identified parasites in the study. *S. haematobium* and *E. vermicularis* recorded 5.6% and 3.7% prevalence rate respectively. The observation is worrisome because it falls out of range of the report of Cameron and Padian (1990). The researchers reported that the prevalence of *T. vaginalis* could be as low as 5% in healthy women and as high as 50% in prostitutes. The high prevalence of *T. vaginalis* reported in this study is however higher than the results recorded in other parts of Nigeria. Akinbo *et al* (2017) reported 2.8% among pregnant women in Benin city, Ijeoma *et al* (2018) observed 40% prevalence among adult in Oru-East local government area of Imo state, Ijasan *et al* (2018) recorded 10% and 8.1% prevalence among HIV-positive and HIV-negative pregnant women in Lagos respectively while Olusola *et al* (2010) recorded 20% among pregnant women in Abeokuta. The result was also higher than 16% reported among women of reproductive age in Brazil (von Glehn *et al.*, 2017) and the 33.3% recorded among Egyptian women by Mahmoud *et al* (2015) and the 34.0% recorded in Nairobi (Mirza *et al.*, 1983).

The variation in observed prevalence could be attributed to the setting (Bowden & Garnett, 2000), the location (Akinbo *et al.*, 2017), socio-economic practices in the study area, laboratory investigative method used, type of specimen and numbers of sexual partners (Fernando *et al.*, 2012; Hussein *et al.* 2015; Hamdy and Hamdy, 2018).

In this study, more females 86/97(88.7%) were infected than males 11/97(11.3%). This may be attributed to the large numbers of females examined and the presence of risk factors in females that promote the striving of the parasite. Factors such as menstruation, sexual activities, pregnancy, vaginal pH value of between 5.0 to 6.0 tend to influence the growth of the parasite (Jirovec & Petru, 1968; Cameron & Garnett, 2000; Agboola, 2006; Swygard *et al.*, 2004).

Although our study indicated that statistically, age was not a risk factor in the prevalence of urinary tract parasitic infections, the parasites were numerically high in students within the age range of 21-25years old. This could be attributed to the fact that this age group are more sexually active (Akinbo *et al.*, 2017). The result is in consonance with the reports of other investigators (Sobel, 2005; Olusola *et al.*, 2010; Mahmoud *et al.*, 2015) but at variance with the record of Akinbo *et al* (2017) that reported a high prevalence in age group of 30-40years old.

Again, the result of our study revealed that the prevalence of urinary tract parasitic infections and specifically *T. vaginalis* was high in single women than married ones. This result conforms with previous studies (DeWaaaji *et al.*, 2017, Okpara *et al.*, 2009, Akinbo *et al.*, 2017). This trend may be as a result of the fact that single women tend to have more than one sexual partner (Usanga *et al.*, 2009, Akinbo *et al.*, 2017; De Waaaji *et al.*, 2017).

Although there was no statistical relationship between nature of toilet and prevalence of urinary tract parasitic infections, women that used bucket system at home had the highest prevalence (54.2%). This result agrees with the findings of Akinbo *et al* (2017).

*E. vermicularis* and *S. haematobium* had a prevalence rate of 3.7% and 5.6% respectively. *E. vermicularis* has been reportedly found in the urine of a female patient by Choudhury *et al* (2017). The parasite is a common gastrointestinal parasite, however, in women, it may be uncommonly found in the vagina, vulva, fallopian tube and uterus. The parasite has been implicated in causation of urinary tract infection especially in regions of high prevalence or endemicity (Ok *et al.*, 1999). The parasite may migrate by ectopic movement from the rectum to the urinary tract.

Urinary infection caused by *E. vermicularis* may be asymptomatic, however, symptoms such as frequent urination, pruritis and itching have been observed in infected females (Choudhury *et al.*, 2017). The presence of the eggs of *S. haematobium* have been reported by many researchers, and the parasite has been mentioned as one of the major causes of urinary bladder carcinoma (Koonrungsesomboon, 2015; Mor *et al.*, 2016).

## Conclusion

The high prevalence rate of urinary tract parasitic infections particularly *T. vaginalis* among the study group is a serious source of concern, and calls for more attention by relevant agencies. Health education, regular screening and treatment of infected persons will reduce the trend.

## Acknowledgement

We appreciate the assistance rendered by the management and staff of Rivers state college of Health Science and Tecnology, Port Harcourt.

## References

1. Agboola A (2006). Vaginal discharge. In: The Textbook of Obstetrics and Gynecology for Medical Students. 2nd ed. Ibadan: Heinemann Educational Books; 2006. p. 548.
2. Akinbo FO, Mokobia CN, Ande AB (2017). Prevalence of trichomoniasis among pregnant women in Benin City. *Sahel Med J* 20:67-71.
3. Bowden FJ, Garnett GP (2000). *Trichomonas vaginalis* epidemiology: Parameterising and analysing a model of treatment interventions. *Sex Transm Infect* 76:248-56
4. Cameron DW, Padian NS (1990). Sexual transmission of HIV and the epidemiology of other sexually transmitted diseases. *AIDS* 4 Suppl 1:S99-103.
5. Choudhury S, Kumar, B, Pal, K.D (2017). *Enterobius vermicularis* infestation of urinary tract leading to recurrent urinary tract infection. *Trop Parasitol* 7(2):119-121
6. Cotch MF, Pastorek JG, Nugent RP, Hillier SL, Gibbs RS, Martin DH, Eschenbach DA, Edelman R, Carey JC, Regan JA, Krohn MA, Klebanoff MA, Rao AV, Rhoads GG (1997) *Trichomonas vaginalis* associated with low birth weight and preterm delivery. The Vaginal Infections and Prematurity Study Group. *Sex Transm Dis* 24: 353-360.
7. de Waaij DJ, Dubbink JH, Ouburg S, Peter RPH, Morrel SA(2017). Prevalence of *Trichomonas vaginalis* infection and protozoan load in South African women: a cross-sectional study. *BMJ Open*;7:e016959. doi:10.1136/ bmjopen-2017-016959
8. Fernando SD, Herath S, Rodrigo C, Rajapakse L (2012). Clinical features and sociodemographic factors affecting *T. vaginalis* infection in women attending a central sexually transmitted diseases clinic in Sri Lanka. *Indian J Sex Transm Dis.* 33, 1:25-31.
9. Fichorova RN (2009). Impact of *T. vaginalis* infection on innate immune responses and reproductive outcome. *J Reprod Immunol* 83:185-9.
10. Fouts AC, Kraus SJ (1980). *Trichomonas vaginalis*: Reevaluation of its clinical presentation and laboratory diagnosis. *J Infect Dis* 141:137-43
11. Gookin JL, Birkenheuer AJ, St John V, Spector M, Levy MG (2005). Molecular characterization of trichomonads from feces of dogs with diarrhea. *J Parasitol* 91:939-43.
11. Hamdy DA, Hamdy HG (2018). Prevalence, Sociodemographic Factors And Clinical Criteria Of *Trichomonas Vaginalis* Infection Among Symptomatic Women In Beni-Suef Governorate, Egypt J. Egypt. Soc. Parasitol. (JESP), 48(1): 109 – 117


12. Hirt RP, de Miguel N, Nakjang S, Dessi D, Liu YC, Diaz N, Rappelli P, Acosta-Serrano A, Fiori PL, Mottram JC (2011) *Trichomonas vaginalis* pathobiology new insights from the genome sequence. *Adv Parasitol* 77: 87-140.
13. Hussein AH, Saleh MH, Nagaty IM, Ghieth KA, El-Azab NA (2015). Prevalence, Clinical Criteria and Sociodemographic Predictors of *Trichomonas vaginalis* infection in suspected Egyptian women, using direct diagnostic techniques. *Iran J. Parasitol.* 10, 3:432-40
14. Ijasan O, Okunade KS, Oluwole AA (2018). The prevalence and risk factors for *Trichomonas vaginalis* infection amongst human immunodeficiency virus-infected pregnant women attending the antenatal clinics of a university teaching hospital in Lagos, South-Western, Nigeria. *Niger Postgrad Med J* 25:21-6.
15. Ijeoma AO, Amara VN, Emmanuel IO (2018). Prevalence of Trichomoniasis among Adults in Oru-East L.G.A, Imo State, Nigeria. *Arch Clin Microbiol.* 9(2):79
16. Jatau ED, Olonitola OS, Olayinka AT (2006). Prevalence of trichomonas infection among women attending antenatal clinics in Zaria, Nigeria. *Ann Afr Med* 5:178-81.
17. Jírovec O, Petru M (1968). *Trichomonas vaginalis* and trichomoniasis. *Adv Parasitol* 6:117-88.
18. Kissinger P (2013). Trichomoniasis AA. and HIV interactions: a review. *Sex Transm Infect* 89:426–33.
19. Koonrungsesomboon N, Wadagni AC, Mbanefo EC (2015). Molecular markers and Schistosoma-associated bladder carcinoma: A systematic review and meta-analysis. *Cancer Epidemiol* 39(4):487-496.
20. Mahmoud A, Sherif NA, Abdella R, El-Genedy, AR, El Kateb AY, Askalani ANA (2015). Prevalence of *Trichomonas vaginalis* infection among Egyptian women using culture and latex agglutination: A cross sectional study. *BMC Women's Health* 15:7.
21. Mann JR, McDermott S, Gill T (2010) Sexually transmitted infection is associated with increased risk of preterm birth in South Carolina women insured by Medicaid. *J Matern Neonatal Med* 23: 563-568.
22. Mirza NB, Nsanze H, D'Costa LJ, Piot P (1983). Microbiology of vaginal discharge in Nairobi, Kenya. *Br J Vener Dis* 59:186-8.
23. Moodley P, Wilkinson D, Connolly C, Moodley J, Sturm AW (2002) *Trichomonas vaginalis* is associated with pelvic inflammatory disease in women infected with human immunodeficiency virus. *Clin Infect Dis* 34: 519-522.
24. Mor N, Tekdoğan UY, Bağcıoğlu M (2016). Parasitic Diseases of Urinary Tract. *Middle Black Sea Journal of Health Science* 2(3):11-1
25. Nester EW, Anderson DG, Roberts CE, Pearsall NN, Nester MT (2001). Human perspective microbiology McGraw-hillinc. Company 3: 657658.
26. Ok UZ, Ertan P, Limoncu E, Ece A, Ozbakkaloglu B(1999). Relationship between pinworm and urinary tract infections in young girls. *APMIS* 107:474–6.
27. Okpara K, Udoidung N, Ating I, Bassey E, Okon O, Nwabueze A. (2009). Risk factors for vaginal Trichomoniasis among women in Uyo, Nigeria. *Internet J Health* 9:1528-8315
28. Olusola O, Taiwo BO, Babatunde DO, Olawunmi SR, Oloyede SB, Adegboyega A (2010). Prevalence of *Trichomonas vaginalis* Infection among Pregnant Women in Abeokuta, Nigeria. *Sierra Leone Journal of Biomedical Research*, 2 (2): 82-86.
29. Patel EU, Gaydos CA, Packman ZR, Quinn TC, Tobian AAR (2018). Prevalence and Correlates of *Trichomonas vaginalis* Infection Among Men and Women in the United States. *Clin Infect Dis.* 67(2): 211-217 doi: 10.1093/cid/ciy079
30. Quinlivan EB, Patel SN, Grodensky CA, Golin CE, Tien HC, Hobbs MM (2012). Modeling the impact of *Trichomonas vaginalis* infection on HIV transmission in HIV-infected individuals in medical care. *Sex Transm Dis* 39:671-7.

31. Ryu JS, Roh J, Lim YS, Seo MY, Choi Y (2015) The secretory products of *Trichomonas vaginalis* decrease fertilizing capacity of mice sperm in vitro. *Asian J Androl* 17: 319.
32. Sam-Wobo SO, Ajao OK, Adeleke MO, Ekpo UF (2012). Trichomoniasis among Antenatal attendees in a tertiary health facility, Abeokuta, Nigeria. *Mun Ent Zool* 7:380-4
33. Sobel JD (2005). What's new in bacterial vaginosis and trichomoniasis? *Infect Dis Clin North Am* 19:387-406.
34. Swygard H, Sena AC, Hobbs MM, Gohen MS (2004) Trichomoniasis clinical manifestations diagnosis and management. *Sexually Transmitted Infections* 80: 91-95.
35. Toth MD (2013) Trichomoniasis and Infertility. *Journal of infectious diseases* 19: 68-70.
36. Upcroft P, Upcroft JA (2001). Drug targets and mechanisms of resistance in the anaerobic protozoa. *Clin Microbiol Rev* 14:150-64.
37. Usanga V, Abia-Basse L, Inyang-Etoh P, Udoh S, Ani F, Archibong E (2009). *Trichomonas vaginalis* infection among pregnant women in Calabar, Cross River State, Nigeria. *Int J Gynecol Obstet* 14:1-7.
38. Van der Pol B. (2007). *Trichomonas vaginalis* infection: the most prevalent nonviral sexually transmitted infection receives the least public health attention. *Clin Infect Dis* 44:23-5.
39. von Glehn M, Ferreira Sá LCE, da Silva HDF, Machado ER (2017). Prevalence of *Trichomonas vaginalis* in women of reproductive age at a family health clinic. *J Infect Dev Ctries* 11(3):269-276. doi:10.3855/jidc.8143
40. WHO (2001). Global prevalence and incidence of selected curable sexually transmitted infections. Overview and Estimate; Geneva 12: 156.
41. WHO (2008). DoRHaR. Global incidence and prevalence of selected curable sexually transmitted infections - 2008. 2012. Report No.
42. WHO, 2012: Global incidence and prevalence of selected curable sexually transmitted infections: Available at <http://www.who.int/reproductivehealth/publications/rtis/stisestimates/en>.






**Plate 1:** *Trichomonas .vaginalis*



**Plate 2:** Egg of *S.haematobium*



**Plate 3:** Egg of *Enteriobius Vermicularis*